

Anti-CD278/ICOS (C398.4A)-169Tm

Catalog number, package size: 3169030B, 100 tests
 3169030C, 25 tests

Clone: C398.4A

Other Names: Inducible co-stimulatory molecule, H4

Isotype: Armenian Hamster IgG

Reactivity: Human, Rhesus, Rat, Mouse, Porcine

Tag: 169Tm

Formulation: Antibody stabilizer with 0.05% sodium azide

Storage: Store at 2–8 °C. Do not freeze.

Application: CyTOF® suspension mass cytometry


Technical Information

Description: CD278, also known as ICOS, is a 50–60 kDa homodimeric membrane glycoprotein and a member of the CD28 family reacting with the Inducible Co-Stimulatory Ligand (ICOSL) molecule. It is highly expressed on activated T cells. It is the receptor for B7-related protein 1 (B7RP-1). Like CD28, ICOS is a co-stimulatory signal for T cell activation and proliferation and cytokine production. It is not expressed on resting or activated B cells, monocytes, NK cells, granulocytes, dendritic cells, or platelets. Unlike the constitutively expressed CD28, ICOS expression is de novo. It has been suggested that ICOS may play an important role in IL-10 production. In the presence of IL-10, purified recombinant human ICOS significantly increased in vitro B cell growth stimulated by pokeweed mitogen (PWM) and enhanced production of IgG.

Application: The metal-tagged antibody is designed and formulated for the application of suspension mass cytometry using the Fluidigm CyTOF suspension systems on healthy human PBMC.

Validation: Each lot of conjugated antibody is quality control-tested by CyTOF suspension mass cytometry analysis of stained cells using appropriate positive and negative cell staining and/or activation controls.

Recommended use: Use 1 µL for up to 3×10^6 live cells in 100 µL staining volume. We recommend titrating the antibody for optimal performance for each of the desired applications. Centrifuge the stock antibody at $12,000 \times g$ for 5 min to sediment antibody aggregates. Fixation is typically used in intracellular staining protocols or in barcoding with the Cell-ID™ 20-Plex Pd Barcoding Kit (PN 201060). However, fixing before antibody staining can affect epitope structure and antibody binding, with the impact varying on the type and concentration of fixative and the protocol used. It is therefore important to perform a small, preliminary antibody staining experiment, with and without fixation, using non-critical samples.



PHA-stimulated human PBMC were stained with anti-CD4 (RPA-T4)-145Nd and anti-ICOS (C398.4A)-169Tm antibodies. The plot shown on the left is gated on total NK cells (CD45+CD3-CD20-CD14-CD56+), and the plot shown on the right is gated on total T cells (CD45+CD20-CD14-CD3+).

Applicable Protocols

Before using this product, refer to the instructions in the Maxpar® Cell Surface Staining with Fresh Fix Protocol (400276).

References

Bandura, D.R. et al. "Mass cytometry: technique for real time single cell multitarget immunoassay based on inductively coupled plasma time-of-flight mass spectrometry." *Analytical Chemistry* 81 (2009): 6,813–22.

Ornatsky, O.I. et al. "Highly multiparametric analysis by mass cytometry." *Journal of Immunological Methods* 361 (2010): 1–20.

Bengsch, B. et al. "Epigenomic-guided mass cytometry profiling reveals disease-specific features of exhausted CD8 T cells." *Immunity* 48 (2018): 1,029–45.

Safety

Use standard laboratory safety protocols. Read and understand the safety data sheets (SDSs) before handling chemicals. To obtain SDSs, go to fluidigm.com and search for 3000000X.

For technical support visit techsupport.fluidigm.com. | For general support visit fluidigm.com/support.

For Research Use Only. Not for use in diagnostic procedures.

Information in this publication is subject to change without notice. **Patent and license information:** fluidigm.com/legal/notices. **Limited Use Label License:** The purchase of this Fluidigm Instrument and/or Consumable product conveys to the purchaser the limited, non-transferable right to use with only Fluidigm Consumables and/or Instruments respectively except as approved in writing by Fluidigm. **Trademarks:** Fluidigm, the Fluidigm logo, Cell-ID, CyTOF, and Maxpar are trademarks and/or registered trademarks of Fluidigm Corporation in the United States and/or other countries. © 2020 Fluidigm Corporation. All rights reserved. 07/30/2020