

Anti-Human BRCA1-172Yb

Pathologist-Verified Clone for Imaging Mass Cytometry™

Catalog: 3172030D

Package size and concentration: 25 µg, 0.5 mg/mL

Storage: Store at 4 °C. Do not freeze.

Reactivity: Human

Clone: MS110

Isotype: Mouse IgG1

Formulation: Antibody stabilizer with 0.05% sodium azide

Application: IMC-Paraffin

Technical Information


Application: The metal-tagged antibody is designed and formulated for the application of Imaging Mass Cytometry (IMC™) using the Fluidigm Hyperion™ Imaging System on formalin-fixed, paraffin-embedded (FFPE) tissue sections.

Quality control: Each lot of conjugated antibody is quality control-tested by Imaging Mass Cytometry on tissue sections.

Recommended concentration: For optimal performance it is recommended that the antibody be titrated for the desired application. Suggested initial dilution range:
IMC-Paraffin: 1:25 to 1:100

Description

BRCA1 (breast cancer type 1 susceptibility protein) is an important member of the DNA repair pathway, and it functions as a tumor suppressor. It interacts with a wide range of proteins involved in the detection of damaged DNA and activation of appropriate repair pathways, including the Mre11-Rad50-NBS1 (MRN) complex, which is responsible for homologous recombination in the repair of DNA double-stranded breaks. BRCA1 also plays a role in cell cycle regulation, where decreased expression of BRCA1 leads to cell cycle arrest through p53 and p21 genes. BRCA1 and BRCA2 are frequently mutated in cases of hereditary breast and ovarian cancer.



Human breast carcinoma (FFPE) stained with 172Yb-anti-BRCA1 (MS110) at a dilution of 1:50 (red pseudocolor) and iridium DNA intercalator (blue pseudocolor). Heat-mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. Scale bar size = 100 µm.

References

Chang, Q. et al. "Staining of frozen and formalin-fixed, paraffin-embedded tissues with metal-labeled antibodies for imaging mass cytometry analysis." *Current Protocols in Cytometry* 82 (2017): 12.47.1–12.47.8.

Giesen, C. et al. "Highly multiplexed imaging of tumor tissues with subcellular resolution by mass cytometry." *Nature Methods* 11 (2014): 417–22.

For technical support visit <http://techsupport.fluidigm.com>. | For general support visit www.fluidigm.com/support.

For Research Use Only. Not for use in diagnostic procedures.

Information in this publication is subject to change without notice. **Safety data sheet information:** www.fluidigm.com/sds. **Patent and license information:** www.fluidigm.com/legalnotices. **Limited Use Label License:** The purchase of this product conveys to the purchaser the limited, non-transferable right to use the purchased consumable or reagent only with Fluidigm Instruments and Systems. **EU's WEEE directive information:** www.fluidigm.com/compliance. Fluidigm, the Fluidigm logo, Hyperion, Imaging Mass Cytometry, and IMC are trademarks or registered trademarks of Fluidigm Corporation in the United States and/or other countries. © 2017 Fluidigm Corporation. All rights reserved. 10/2017